



1 August 1994

PHYSICS LETTERS A

An approximate renormalization for the break-up of invariant tori with three frequencies

R.S. MacKay¹, J.D. Meiss², J. Stark

Centre for Nonlinear Dynamics and Applications, University College London, London WC1E 6BT, UK

Received 8 April 1994, revised manuscript received 21 April 1994, accepted for publication 17 May 1994

Communicated by A.R. Bishop

Abstract

Renormalization theory provides a description of the destruction of invariant tori for Hamiltonian systems of $1\frac{1}{2}$ or 2 degrees

dent resonances for ω , then $\omega = p$ where p is integral (remember the length of ω is unimportant). A frequency ω is *Diophantine* if there is a $K \neq 0$ and $\tau > 2$

tersection of each pair of resonances defines rational frequencies $p_1 = [1, 0, 0]$, $p_2 = [0, 1, 0]$, $p_3 = [0, 0, 1]$. The frequencies p_i also delineate the cone: it is the

such that $\forall m \in \mathbb{Z}^3 \setminus 0, |m \cdot \omega| / |\omega| > K / |m|^\tau$.

When $A=B=C=0$, the momenta (u, v) are constant in time and every orbit lies on a three torus. If $\omega(u, v)$ is incommensurate, the orbit densely covers the torus. If ω is Diophantine, then the KAM theorem implies that there is a torus with this frequency for small values of the amplitudes. We are interested

convex hull of the three vectors. We denote the cone by either of the matrices

$$M = \begin{pmatrix} m_1 \\ m_2 \\ m_3 \end{pmatrix}, \quad P = (p_1, p_2, p_3).$$

We assume ω is inside the cone, i.e. $\omega_i \geq 0$.

1994-03-15 10:00 AM

[19] R.S. MacKay and J. Stark, *Phys Lett A* 138 (1989) 113

Physica D 49 (1991) 387