

1. (20 pts) Parts (a) and (b) are not related.

- (a) For  $f(x) = \frac{1}{x^2}$  and  $g(x) = \frac{1}{x+2}$ , identify the composite function  $(f \circ g)(x)$  and its domain. Express the domain in interval form.

**Solution:**


$$(f \circ g)(x) = f(g(x)) = f\left(\frac{1}{x+2}\right) = \frac{1}{\left(\frac{1}{x+2}\right)^2} = (x+2)^2 = \boxed{x+2}$$

The domain of  $f \circ g$  is the set of all  $x$  in the domain of  $g$  such that  $g(x)$  is in the domain of  $f$ .

$$\text{Domain of } g: \left( x+2 > 0 \right) \implies x > -2$$


For each  $x$  in the interval  $(-2; \infty)$ ,  $g(x)$  is in the domain of  $f$  (since  $g(x) \neq 0$  for all  $x$ ).

- (b) The graph below depicts a function of the form  $y = h(x) = a \sin (bx) + c$ . Determine the values of  $a$ ,  $b$ , and  $c$ . (*Hint: Consider the transformations from the graph of  $y = \sin x$  to the given graph.*)



**Solution:**

Begin with the graph of the relevant base curve,  $y = \sin x$ :



The profile of the given curve over the interval  $[0; \pi]$  is the same as the profile of the  $y = \sin x$  curve over the interval  $[0; 3\pi]$ . Therefore, the given curve has experienced a horizontal compression of a factor of 3 with respect to the  $y = \sin x$  curve, which implies that  $b = 3$

The vertical difference between the given curve's maximum and minimum values is  $1 - (-3) = 4$ , while the vertical difference between the  $y = \sin x$  curve's maximum and minimum values is  $1 - (-1) = 2$ . Therefore, the given curve has experienced a vertical expansion of a factor of 2 with respect to the  $y = \sin x$  curve, which implies that  $a = 2$

The vertical center of the given curve is  $y = -1$  while the vertical center of the  $y = \sin x$  curve is  $y = 0$ . Therefore, the given curve has experienced a downward vertical shift of 1 unit with respect to the  $y = \sin x$  curve, which implies that  $c = -1$

Therefore, the function depicted in the given graph is  $y = 2 \sin (3x) - 1$

2. (30 pts) Evaluate the following limits. Support your answers by stating theorems, definitions, or other key properties that are used.

(a)  $\lim_{x \neq 0} \frac{\tan x \sin(2x)}{x^2}$

**Solution:** Key property:  $\lim_{x \neq 0} \frac{\sin x}{x} = 1$

$$\begin{aligned} \lim_{x \neq 0} \frac{\tan x \sin(2x)}{x^2} &= \lim_{x \neq 0} \frac{\tan x}{x} \cdot \frac{\sin(2x)}{x} \\ &= \lim_{x \neq 0} \frac{\sin x}{x \cos x} \cdot \frac{2 \sin(2x)}{2x} \\ &= \lim_{x \neq 0} \frac{\sin x}{x} \cdot \frac{1}{\cos x} \cdot \frac{2 \sin(2x)}{2x} \\ &= \lim_{x \neq 0} \frac{\sin x}{x} \cdot \lim_{x \neq 0} \frac{2}{\cos x} \cdot \lim_{x \neq 0} \frac{\sin(2x)}{2x} \\ &= [1] \cdot \frac{2}{1} \end{aligned}$$

$$(b) \lim_{x \rightarrow 9} \frac{\sqrt{x-5} - 2}{x-9}$$

**Solution:**

Begin by multiplying the numerator and the denominator by the conjugate of the original numerator expression.

$$\lim_{x \rightarrow 9} \frac{\sqrt{x-5} - 2}{x-9} = \lim_{x \rightarrow 9} \frac{\sqrt{x-5} - 2}{x-9} \cdot \frac{\sqrt{x-5} + 2}{\sqrt{x-5} + 2}$$

3. (30 pts) Consider the rational function  $r(x) = \frac{x^2 - 5x + 4}{2x^2 - 8x + 6}$ .

(a) Identify all values of  $x$  at which  $r(x)$  is discontinuous. At each such  $x$  value, explain why the function is discontinuous there.

**Solution:**

$$r(x) = \frac{x^2 - 5x + 4}{2x^2 - 8x + 6} = \frac{(x - 1)(x - 4)}{2(x - 1)(x - 3)}$$

Since  $r(x)$  is a rational function, it is continuous at all  $x$  in its domain.

Therefore,  $r(x)$  is discontinuous only at  $x = 1; 3$

(b) Identify the type of discontinuity associated with each  $x$  value identified in part (a). Support those classifications by evaluating the appropriate limits.

**Solution:**

$$r(x) = \frac{(x - 1)(x - 4)}{2(x - 1)(x - 3)} = \frac{(x - 4)}{2(x - 3)}, \quad x \neq 1; 3$$

$$\lim_{x \rightarrow 1} r(x) = \lim_{x \rightarrow 1} \frac{x - 4}{2(x - 3)} = \frac{1 - 4}{(2)(1 - 3)} = \frac{-3}{-4} = \frac{3}{4}$$

Since the two-sided limit is finite, there is a removable discontinuity at  $x = 1$

$$\lim_{x \rightarrow 3^-} r(x) = \lim_{x \rightarrow 3^-} \frac{x - 4}{2(x - 3)} = \frac{1}{(2)(0^-)} = -\infty$$

$$\lim_{x \rightarrow 3^+} r(x) = \lim_{x \rightarrow 3^+} \frac{x - 4}{2(x - 3)} = \frac{1}{(2)(0^+)} = \infty$$

Since at least one of the two preceding one-sided limits is infinite, there is an infinite discontinuity at  $x = 3$

- (c) Find the equation of each vertical asymptote of  $y = r(x)$ , if any exist. Support your answer in terms of the limits you evaluated in part (b).

**Solution:**

The finite value of  $\lim_{x \rightarrow 1} r(x) = \frac{3}{4}$  determined in part (b) indicates that there is no vertical asymptote at  $x = 1$ .

The infinite limits  $\lim_{x \rightarrow 3} r(x) = 1$  and  $\lim_{x \rightarrow 3^+} r(x) = 1$  were determined in part (b). Either one of those limits being infinite is sufficient to conclude that the line  $x = 3$  is a vertical asymptote of the curve  $y = r(x)$ .

- (d) Find the equation of each horizontal asymptote of  $y = r(x)$ , if any exist. Support your answer by evaluating the appropriate limits.

**Solution:**

$$\begin{aligned}\lim_{x \rightarrow 1} r(x) &= \lim_{x \rightarrow 1} \frac{x^2 - 5x + 4}{2x^2 - 8x + 6} = \lim_{x \rightarrow 1} \frac{x^2 - 5x + 4}{2x^2 - 8x + 6} \cdot \frac{1-x^2}{1-x^2} \\ &= \lim_{x \rightarrow 1} \frac{1 - 5 + 4 - x^2}{2 - 8 + 6 - x^2} = \frac{1 - 0 + 0}{2 - 0 + 0} = \frac{1}{2}\end{aligned}$$

Therefore, the equation of the only horizontal asymptote is  $y = \frac{1}{2}$

4. (20 pts) Parts (a) and (b) are not related.

(a) For what value of  $b$  is the following function  $u(x)$  continuous at  $x = 3$ ? Support your answer using the definition of continuity, which includes evaluating the appropriate limits.

$$u(x) = \begin{cases} \frac{x^2 - 9}{x - 3} & ; x < 3 \\ 5x + b & ; x \geq 3 \end{cases}$$

**Solution:**

By the definition of continuity,  $u(x)$  is continuous at  $x = 3$  if  $\lim_{x \rightarrow 3^-} u(x) = \lim_{x \rightarrow 3^+} u(x) = u(3)$ .

$$\lim_{x \rightarrow 3^-} u(x) = \lim_{x \rightarrow 3^-} \frac{x^2 - 9}{x - 3} = \lim_{x \rightarrow 3^-} \frac{(x - 3)(x + 3)}{x - 3} = \lim_{x \rightarrow 3^-} (x + 3) = 3 + 3 = 6$$

$$\lim_{x \rightarrow 3^+} u(x) = \lim_{x \rightarrow 3^+} (5x + b) = (5)(3) + b = 15 + b$$

$$u(3) = (5)(3) + b = 15 + b$$

Therefore,  $u(x)$  is continuous at  $x = 3$  if  $6 = 15 + b$ , which occurs when  $b = -9$

(b) The Intermediate Value Theorem can **NOT** be used to guarantee that  $v(x) = \frac{2}{x} + \frac{1}{x+2} = 0$  for a value of  $x$  on the interval  $(-1; 2)$ . Explain which condition for applying the theorem is not satisfied in this case.

**Solution:**

The Intermediate Value Theorem cannot be applied in this case because  $v(0)$  is undefined, which means that

$v(x)$  is not continuous on the interval  $[-1; 2]$

The continuity of  $v(x)$  on  $[-1; 2]$  is one of the hypotheses for applying the IVT to the given function on the given interval.

(Note that  $v(-1) = -1$  and  $v(2) = 3$  together indicate that the other IVT hypothesis does hold)