smaller solitary wave increases in amplitude and experiences a negative position shift (Zabusky & Kruskal 1965). The two solitary waves then emerge from the interaction

FIGURE 1. (Colour online) Diagram of the uid conduit experimental system.

Using careful numerical simulations, we nd that although the conduit equation



FIGURE 4. (Colour online) Data from interaction experiments. (*a*) Unscaled data showing the solitary wave profiles from a typical interaction. (*b*) Examples of the three interaction types for scaled data used in classifications. From top to bottom, $.a_{lead}; a_{trail}/$ are .1:27; 3:45/, .1:53; 9:97/, and .1:14; 10:01/, respectively. See the online version of the paper for movies of these experiments.

differences and residuals are beyond the sensitivity of our experimental capabilities, which suggests that these conduit solitary waves are approximately solitons, at least in a physical sense. Hence we term them 'physical solitons'.

3.4. *Experimental observation of the three types of interaction*

In figure 3, we plot the results of twenty-seven solitary wave interaction classification experiments. The three distinct types predicted by numerical simulations of the conduit equation (2.5) are readily observable in the full physical system, and their dependence on a_{lead} and a_{trail} is in excellent agreement with the phase diagram. Example images of an unscaled interaction experiment and then scaled data used for classification are given in figure 4. While it is sometimes difficult to distinguish between the regimes in the unscaled data, scaling the data recovers the aspect ratio of the nondimensional coordinate system from the numerical simulations and allows for proper determination. Typical examples of the three interaction types are shown. Movies in both unscaled and scaled formats are available with the online version of the paper at http://dx.doi.org/10.1017/jfm.2014.273.

Regarding the agreement with numerical simulations, up to the 2% error in measuring conduit diameters, the data all lie in the appropriate regions of the phase diagram. While this agreement is compelling on its own, it is also possible to compare

with the approximate model breakdown criteria identified in Lowman & Hoefer (2013*a*). The use of miscible fluids meets the negligible surface tension criterion, and the contributions due to the outer wall are small because the nondimensional radius of the outer wall satisfies R_{wall} 76 $^{1=2}$ D 10:3. The Reynolds number criterion *Re* D 0:049 ^{1} D 106 for neglecting inertial effects is also satisfied. Lastly, the breakdown of the multiple scales assumption occurs when the condition, *A* 1=8 13:3, is violated for large amplitude solitary waves (Lowman & Hoefer 2013*a*). We note that two of the trials lie beyond this point, though they still fall in the appropriate classification region. There exists an independent condition for model validity based on the introduction of inertial effects for sufficiently large amplitude solitary waves, measured by an effective solitary wave Reynolds number (Helfrich & Whitehead 1990), but we find the multiple scales condition to be more restrictive in our case. Our results suggest remarkably robust concurrence between the reduced, approximate interfacial equation and the full two-fluid system.

4. Summary and conclusions

The qualitative characterization of large amplitude, pairwise solitary wave interactions in viscous fluid conduits has been shown to permit geometric classification according to the three Lax categories for KdV. Unlike the weakly nonlinear regime, however, the expected interaction type depends on the wave amplitudes, rather than only their ratio, and the mixed unimodal/bimodal interaction type is a more robust, readily observable feature than for surface water waves.

The long time, large amplitude validity of the conduit equation (2.5) and its analytical tractability make this two viscous fluid setting an ideal one for the study of nonlinear dispersive waves. That nonlinear dispersive waves occur at all in a fully viscous setting is a nontrivial observation, but that the reduced equation captures the geometry of interacting solitary waves suggests the interfacial dynamics of viscous fluid conduits are, as predicted, approximately one-dimensional and dissipationless at the time scales under consideration. Moreover, the absence of dispersive radiation in the experiments implies that, while the conduit equation is not completely integrable, its solitary waves practically interact elastically. These results encourage future experimental studies on nonlinear coherent structures, such as rarefaction waves, slowly modulated wavetrains (dispersive shock waves) and their interactions.

Acknowledgements

We thank K. Daniels, H. Tran, and the NCSU Department of Mathematics for laboratory support. This work was funded by NSF Grant Nos. DGE-1252376, DMS-1008793, and CAREER DMS-1255422.

Supplementary movies

Supplementary movies are available at http://dx.doi.org/10.1017/jfm.2014.273.

REFERENCES

ABLOWITZ, M. J. & HAUT, T. S. 2010 Asymptotic expansions for solitary gravity–capillary waves in two and three dimensions. *J. Phys.* A **43** (43), 434005.

BARCILON, V. & RICHTER, F. M. 1986 Nonlinear waves in compacting media. *J. Fluid Mech.* **164**, 429–448.

BENJAMIN, T. B. 1967 Internal waves of permanent form in fluids of great depth. *J. Fluid Mech.* **29** (3), 559–592.