

Preliminary Exam
 Partial Differential Equations
 9:00 AM - 12:00 PM, Jan. 11, 2024
 Newton Lab, ECCR 257

Student ID (do NOT write your name):

#	possible	score
1	25	
2	25	
3	25	
4	25	
5	25	
Total	100	

There are five problems. **Solve four of the five problems.**
 Each problem is worth 25 points.

A sheet of convenient formulae is provided.

1. **Method of characteristics.** Consider the inviscid Burger's equation

$$u_t + uu_x = 0 \tag{1}$$

on the domain $\Omega = \mathbb{R} \times \mathbb{R}^+$ with initial conditions

$$u(x, 0) = u_0(x) = \begin{cases} 1, & x \leq 0, \\ 1 - x, & 0 < x < 1, \\ 0, & 1 < x. \end{cases} \tag{2}$$

(a) Find the time and position at which a shock forms.

Solution: The characteristic equations are

$$\frac{dt}{d\tau} = 1, \tag{3}$$

$$\frac{dx}{d\tau} = u, \tag{4}$$

$$\frac{du}{d\tau} = 0, \tag{5}$$

$$\tag{6}$$

which gives, using the initial data $(x, t, u) = (s, 0, u_0(s))$,

$$t = \tau, \tag{7}$$

$$x = ut + s, \tag{8}$$

$$u = u_0(s). \tag{9}$$

Thus, the solution u satisfies the implicit equation $u = u_0(x - ut)$. To find the location of the shock, we differentiate with respect to x and solve for u_x , finding

$$u_x = \frac{u_0'}{1 + u_0 t}. \tag{10}$$

Thus, a characteristic emanating from the initial point $(s, 0)$ has a slope of $u_0(s)$. The characteristics emanating from the initial point $(s, 0)$ are straight lines in the (x, t) plane with slope $u_0(s)$. The characteristics emanating from the initial point $(s, 0)$ are straight lines in the (x, t) plane with slope $u_0(s)$. The characteristics emanating from the initial point $(s, 0)$ are straight lines in the (x, t) plane with slope $u_0(s)$.

we conclude that all characteristics emanating from $(0, 1)$ produce a shock at $t_s = 1$. The position of the shock for the characteristic starting at $x_0 = s \in (-1, 1)$ can be found by setting $t = t_s = 1$ and $u = u_0(s) = 1 - s$ in Eq. (8), which gives $x_s = (1 - s)1 + s = 1$. Therefore the shock forms at $(x_s, t_s) = (1, 1)$.

- (b) Find the subsequent trajectory of the discontinuous shock by applying the Rankine-Hugoniot condition

$$s(t) = \frac{1}{2}(u_-(t) + u_+(t)),$$

where s is the speed of the discontinuity and $u_{\pm}(t) = \lim_{x \rightarrow x_s(t) \pm} u(x, t)$ and $s = \dot{x}_s(t)$.

Solution: Since the Burgers equation can be written as $u_t + (u^2/2)_x = 0$, the Rankine-Hugoniot condition for the position of the shock $x_s(t)$ gives

$$\frac{dx_s}{dt} = \frac{\frac{1}{2}u_+^2 - \frac{1}{2}u_-^2}{u_+ - u_-}, \quad (12)$$

where u_+ and u_- are the values of u to the right and to the left of the shock, respectively. The value to the left corresponds to characteristics emanating from $x_0 < 0$, for which $u = 1$, and the value to the right corresponds to characteristics emanating from $x_0 > 1$, for which $u = 0$ (a rough sketch of the characteristics might be useful here). Thus, $u_+ = 0$ and $u_- = 1$, and we have

$$\frac{dx_s}{dt} = \frac{\frac{1}{2}0 - \frac{1}{2}1}{0 - 1} = \frac{1}{2}. \quad (13)$$

Together with the initial condition $x_s(1) = 1$, we get $x_s(t) = 1 + (t - 1)/2$.

- (c) Sketch the characteristics and the shock in the (x, t) plane.

Solution: A sketch is shown below.

- (d) Find the solution $u(x, t)$.

Solution: The solution satisfies the implicit equation $u = u_0(x_0) = u_0(x - ut)$. When $x_0 < 0$, $u_0 = 1$, and so we have $u = 1$ along the characteristics $x_0 = x - t$ for $x_0 < 0$, provided they haven't met the shock (blue lines in diagram). Similarly, $u_0 = 0$ for $x_0 > 0$, and so $u = 0$ along the characteristics $x_0 = x$ for $x_0 > 0$ (purple lines). Finally, if $0 < x_0 < 1$ we have $u_0 = 1 - x_0$, and so $u = 1 - (x - ut)$, which yields $u = (1 - x)/(1 - t)$ (green lines). Putting everything together, we obtain

3. **Wave Equation.** Consider the following initial-boundary value problem on the domain $D = \{(x, t) : t \in \mathbb{R}^+, x \in \mathbb{R}^+, x > t/\alpha\}$, where $\alpha > 1$:

$$u_{tt} = u_{xx}, \quad x > t/\alpha, \quad t > 0, \quad (15)$$

$$u(x, 0) = \phi(x), \quad x > 0, \quad (16)$$

$$u_t(x, 0) = \psi(x), \quad x > 0, \quad (17)$$

$$u(x, x/\alpha) = f(x), \quad x > 0, \quad (18)$$

with $\phi, \psi, f \in C^2(\mathbb{R}_0^+)$.

(a) Find the solution $u(x, t)$.

Solution: We seek a solution of the form

$$u(x, t) = F(x - t) + G(x + t) \quad (16)$$

(b) Find sufficient conditions on α , β , and f so that the solution is continuous in D .

Solution: We need to ensure continuity across $x = t$, where the two solutions meet. Letting $x = t^+$ and using the fact that the functions involved are continuous we get

lim

Multiplying the PDE by v and integrating over the domain, we have

$$\begin{aligned} 0 &= \int_{B(0,1)} v(\mathbf{x}) \Delta v(\mathbf{x}) \, d\mathbf{x} \\ &= - \int_{B(0,1)} |\nabla v(\mathbf{x})|^2 \, d\mathbf{x} + \int_0^{2\pi} v(1, \theta) v_r(1, \theta) \, d\theta \\ &= - \int_{B(0,1)} |\nabla v(\mathbf{x})|^2 \, d\mathbf{x}, \end{aligned}$$

upon applying integration by parts and the boundary condition. Since the integrand is non-negative definite and the integral is zero, we must have

$$|\nabla v(\mathbf{x})|^2 = 0, \quad \mathbf{x} \in B(0,1) \quad \Rightarrow \quad v(\mathbf{x}) = \text{const}, \quad \mathbf{x} \in B(0,1).$$

Since the average of $v(\mathbf{x})$ on the boundary is zero, $v(\mathbf{x})$ must be identically zero and uniqueness is proven.

- (b) We seek a solution using the method of separation of variables in polar coordinates. Then, eq. (33) becomes

$$\begin{aligned} u_{rr} + \frac{1}{r} u_r + \frac{1}{r^2} u_{\theta\theta} &= 0, \quad r \in (0,1), \quad \theta \in [0,2\pi], \\ u_r(1, \theta) &= g(\theta), \quad \theta \in [0,2\pi]. \end{aligned}$$

Seeking a solution in separated form $u(r, \theta) = f(r)g(\theta)$ implies

$$\begin{aligned} g''(\theta) + \lambda g(\theta) &= 0, \quad \theta \in [0,2\pi], \quad g(0) = g(2\pi), \quad g'(0) = g'(2\pi), \\ f''(r) + \frac{1}{r} f'(r) - \frac{\lambda}{r^2} f(r) &= 0, \quad r \in (0,1), \quad \lim_{r \rightarrow 0} |f(r)| < \infty. \end{aligned}$$

The angular boundary value problem has the trigonometric solutions

$$g_n(\theta) = A_n \cos(n\theta) + B_n \sin(n\theta), \quad n = 0, 1, 2, \dots$$

with the corresponding eigenvalues $\lambda_n = n^2$.

The radial problem exhibits the bounded solutions

$$f_n(r) = r^n.$$

Introduce the series solution

$$u(r, \theta) = A_0 + \sum_{n=1}^{\infty} r^n [A_n \cos(n\theta) + B_n \sin(n\theta)].$$

The coefficients are determined by the boundary conditions

$$u_r(1, \theta) = \sum_{n=1}^{\infty} n [A_n \cos(n\theta) + B_n \sin(n\theta)] = g(\theta), \quad \theta \in [0,2\pi].$$

Multiplying by $\cos(m\theta)$ and integrating from 0 to 2π , we obtain

$$A_m = \frac{1}{m} \int_0^{2\pi} g(\theta) \cos(m\theta) \, d\theta, \quad m = 1, 2, \dots$$

Multiplying by $\sin(m\theta)$ and integrating from 0 to 2π , we obtain

$$B_m = \frac{1}{m} \int_0^{2\pi} g(\theta) \sin(m\theta) \, d\theta, \quad m = 1, 2, \dots,$$

which determines a series representation of the solution. To determine A_0 , we require zero average on the boundary so that $A_0 = 0$.

(c) Inserting the expressions for the coefficients into the series representation, we obtain

$$\begin{aligned}
 u(r, \theta) &= \sum_{n=1}^{\infty} \frac{r^n}{n} \int_0^{2\pi} g(\phi) \cos(n\phi) \cos(n\theta) + \sin(n\phi) \sin(n\theta) \, d\phi \\
 &= \sum_{n=1}^{\infty} \frac{r^n}{n} \int_0^{2\pi} g(\phi) \cos(n(\phi - \theta)) \, d\phi \\
 &= \sum_{n=1}^{\infty} \frac{r^n}{n} N(r, \theta) \, d\phi .
 \end{aligned}$$

where a is a constant and the dot and prime indicate time and space derivatives, respectively. If $a = 0$, the spatial equation gives $X = A + Bx$, which upon evaluation of the boundary conditions leads to $X = 0$. Similarly, if $a > 0$ we get $X = Ae^{\bar{a}x} + Be^{-\bar{a}x}$, leading also to $X = 0$. Therefore, a must be negative and we set $a = -\frac{1}{2}$. We obtain

$$T(t) = T(0) \exp(-\frac{1}{2}t), \quad (42)$$

$$X(x) = A \sin(\frac{1}{2}x) + B \cos(\frac{1}{2}x). \quad (43)$$

Using the boundary conditions $X(0) = X(1) = 0$ we obtain $B = 0$ and $\frac{1}{2} = n$, so we get the modes

$$X_n(x) = \sin(n\pi x), \quad (44)$$

where $\frac{1}{2} = n$ and $n \in \mathbb{N}^+$. Thus, we find

$$\tilde{u}(x, t; s) = \sum_{n=1}^{\infty} A_n e^{-\frac{1}{2}nt} \sin(n\pi x). \quad (45)$$

Using the initial conditions $\tilde{u}(x, t; s) = f(x)e^{-s}$ we get

$$f(x)e^{-s} = \sum_{n=1}^{\infty} A_n e^{-\frac{1}{2}ns} \sin(n\pi x), \quad (46)$$

which implies that $A_n = f_n e^{(\frac{1}{2}n-1)s}$, where f_n is the n th sine Fourier coefficient of $f(x)$. Therefore,

$$\tilde{u}(x, t; s) = \sum_{n=1}^{\infty} f_n e^{(\frac{1}{2}n-1)s} e^{-\frac{1}{2}nt} \sin(n\pi x). \quad (47)$$

and

$$u(x, t) = \int_0^t \tilde{u}(x, t; s) ds = \int_0^t \sum_{n=1}^{\infty} f_n e^{(\frac{1}{2}n-1)s} e^{-\frac{1}{2}nt} \sin(n\pi x) ds \quad (48)$$

$$= \sum_{n=1}^{\infty} f_n e^{-\frac{1}{2}nt} \sin(n\pi x) \int_0^t e^{(\frac{1}{2}n-1)s} ds \quad (49)$$

$$= \sum_{n=1}^{\infty} f_n e^{-\frac{1}{2}nt} \sin(n\pi x) \frac{e^{(\frac{1}{2}n-1)t} - 1}{\frac{1}{2}n - 1} \quad (50)$$

$$= \sum_{n=1}^{\infty} f_n e^{-\frac{1}{2}nt} \sin(n\pi x) \frac{e^{(\frac{1}{2}n-1)t} - 1}{\frac{1}{2}n - 1} \quad (51)$$

$$= \sum_{n=1}^{\infty} f_n \sin(n\pi x) \frac{e^{-t} - e^{-\frac{1}{2}nt}}{\frac{1}{2}n - 1}. \quad (52)$$

(b) Prove that the solution is unique.

Solution: Assume there are two solutions, u_1 and u_2 . Then their difference $w = u_1 - u_2$ satisfies

$$w_t = w_{xx}, \quad 0 < x < 1, \quad t > 0, \quad (53)$$

$$w(x, 0) = 0, \quad 0 < x < 1, \quad (54)$$

$$w(0, t) = w(1, t) = 0 \quad t > 0. \quad (55)$$

Let $T > 0$. By the maximum principle, the maximum of w in the closure of $U_T = [0, 1] \times [0, T)$ must be equal to the maximum of w in its parabolic boundary, $\bar{U}_T - U_T$, which is zero. Therefore $w \leq 0$, or equivalently $u_1 \leq u_2$ in \bar{U}_T . Applying the same argument to $-w$ we conclude that $w = u_1 - u_2 \geq 0$ in \bar{U}_T . Since T was arbitrary, $u_1(x, t) = u_2(x, t)$ for all $t > 0$, $x \in (0, 1)$, so the solution is unique.

END