

Transport through chaos

This content has been downloaded from IOPscience. Please scroll down to see the full text.

1991 Nonlinearity 4 583

(<http://iopscience.iop.org/0951-7715/4/2/017>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 128.138.73.68

This content was downloaded on 05/12/2014 at 21:34

Please note that [terms and conditions apply](#).

Transport through chaos

Robert W Easton†

Department of Mathematics, University of Colorado, Boulder, CO 80309–0426, USA

Received 10 October 1989, in final form 3 April 1990


Accepted by R S Mackay

Abstract. Certain orbits of area preserving maps of the plane appear to wander

and densely fill regions of the plane having positive Lebesgue measure.

determine how ensembles of points are transported. The action principle of MacKay, Meiss and Percival [4] can be used to compute areas of pieces of the grid. Thus knowledge of trellis geometry together with area computations will form the

Figure 2.1. Suppose that the curve μ is a piece of the stable manifold of some




Rearranging the sum gives

$$\int_D dp \wedge dq = \sum_{j=-\infty}^{\infty} \alpha^j [F(b_j) - F(a_j)].$$

In general suppose that D is a disc bounded by alternating segments of stable and unstable manifold. Suppose that the endpoints of these segments are indexed a^0, a^1, \dots, a^{2m} (with $a^{2m} = a_0$) in a counterclockwise order around the boundary of D . Suppose that the segment joining a^0 and a^1 is contained in a stable manifold. Then by the preceding argument


$$\int_D dp \wedge dq = \sum_{j=-\infty}^{\infty} \alpha^j \sum_{k=0}^{m-1} [F(a_j^{2k+1}) - F(a_j^{2k})]. \quad (2.4)$$

This formula expresses the Moser-Moisè-Denisov action principle



Proposition. Discontinuity points of t^+ occur on R -stable manifolds. Similarly, discontinuity points of t^- occur on R -unstable manifolds. Hence the internal trellis of the resonance zone partitions the zone into its exit time decomposition.

Proof. For simplicity the proof will be given for the resonance zone pictured in



- [3] Easton R 1986 Trellises formed by stable and unstable manifolds in the plane *Trans. Am. Math. Soc.* **294** 719–31
- [4] MacKay R, Meiss J and Percival I 1984 Transport in Hamiltonian systems *Physica* **13D** 55–81
- [5] Poincaré H 1892 *Methodes Nouvelles de la Mechanique Celeste* (Paris: Gauthier-Villars) ch 33
- [6] Wiggins S 1990 On the geometry of transport in phase space, 1. Transport in k -degree-of-freedom Hamiltonian systems *Physica* **44D** 471–501
- [7] Rom-Kedar V 1990 Transport rates of a class of two-dimensional maps and flows *Physica* **43D** 229–68
- [8] Easton R 1989 Isolating blocks and epsilon chains for maps *Physica* **39D** 95–110